
MYCOTAXON

Volume 115, pp. 171–173

January–March 2011

DOI: 10.5248/115.171

Stachybotrys subreniformis, new from soil in China

QI-RUI LI & YU-LAN JIANG*

Department of Plant Pathology, Guizhou University, Guiyang, Guizhou, 550025, China

*CORRESPONDENCE TO: yljhsd@163.com

ABSTRACT — A new species of *Stachybotrys* from soil in China, *S. subreniformis*, is illustrated and described. It differs from similar species mainly by producing smaller conidia and shorter conidiophores. The type specimen (dried culture) and living culture are deposited in the Plant Pathology Herbarium of Guizhou University (HGUP).

KEY WORDS — soil fungi, taxonomy, dematiaceous hyphomycetes

Introduction

During an investigation of soil dematiaceous hyphomycetes in Guizhou Province, Southwestern China in 2009, one isolate possessing the typical characters of *Stachybotrys* Corda was obtained from forest soil and found to differ from described species of *Stachybotrys* in conidial and conidiophore morphology. The new taxon is illustrated and described below.

Materials & methods

Fifteen soil samples were collected from Duyun, Guizhou Province. The methods of sample collection, isolation and identification of fungi, and preparation of dried cultures were as previously reported by Jiang & Zhang (2007). Conidia and conidiophores were placed in a drop of 85% lactic acid and examined and photographed using light microscopy. Thirty mature conidia and phialides were measured by using a UB200i microscope (Chongqing UOP Photoelectric Technology Co., China) at 1000× magnification. The fungi are described from cultures grown at 25°C on corn meal agar (CMA).

Taxonomy

Stachybotrys subreniformis Q.R. Li & Y.L. Jiang, sp. nov.

FIG. 1 A–B

MYCOBANK MB 519085

Coloniae effusae, densae, pilosae, griseae vel nigrae. Mycelium superficiale vel immersum. Hyphae hyalinae, ramosae, septatae, laeves, 5–7 μm crassae. Conidiophora macronemata, mononemata, erecta vel flexuosa, ramosa, septatae, hyalinae vel artobrunnea, 48–98 ×




FIG. 1A–B. Conidia and conidiophores of *Stachybotrys subreniformis*. (Bars = 20 μm)

5.0–7.5 μm . Phialides brunnea, obovatae, laeves, 8.0–11.5 \times 4.5–6.0 μm . Conidia sphaerica vel reniformia, aseptata, atrobrunnea, verruculosa, 6.0–9.5 \times 4.5–7.5 μm .

HOLOTYPE: CHINA. GUIZHOU PROVINCE, Duyun, from forest soil, 15 December 2009, Q.R. Li, HGUP₀₉1051.

ETYMOLOGY: in reference to the slightly reniform shape of many of the conidia.

COLONIES on CMA effuse, dense, hairy, grey to black, colourless at first, then becoming dark with production of abundant anamorphic structures. MYCELIUM superficial and immersed: hyphae hyaline, branched, septate, smooth, 5–7 μm wide. CONIDIOPHORES macronematous, mononematous, erect or flexuous, branched, septate, hyaline to dark brown, 48–98 \times 5.0–7.5 μm , tapering toward the apex which bears terminal phialides in a whorl of 2–7 around a central phialide. PHIALIDES brown, obovate, smooth, 8.0–11.5 \times 4.5–6.0 (av. 9.3 \pm 0.9 \times 5.3 \pm 0.4, n = 30) μm , with conspicuous collarettes. CONIDIA aggregated in slimy masses, spherical or slightly reniform, nonseptate, dark brown, verruculose, 6.0–9.5 \times 4.5–7.5 (av. 7.6 \pm 1.0 \times 6.1 \pm 0.8, n = 30) μm .

In the key to *Stachybotrys* species (Pinruan et al. 2004), seven species produced reniform conidia. They are *S. nephrodes* McKenzie (McKenzie 1991), *S. sinuatophora* Matsush. (Matsushima 1971), *S. reniverrucosa* Whitton et al. (Whitton et al. 2001), *S. nephrospora* Hansf. (Hansford 1943), *S. reniformis* Tubaki (Tubaki 1963) and *S. renisporoides* K.G. Karand. et al. (Karandikar et al. 1992) and *S. oenanthes* M.B. Ellis (Ellis 1971). However, only *S. oenanthes* produces ovoid to reniform conidia, which is very close to *S. subreniformis*. Conidia of *S. subreniformis* (6.0–9.5 \times 4.5–7.5 μm) are smaller than those of *S. oenanthes* (9–12 \times 4.5–8 μm). In addition, the conidiophores of *S. subreniformis* (48–98 μm) are shorter than those of *S. oenanthes* (up to 190 μm).

Acknowledgments

The authors are grateful for pre-submission comments and suggestions provided by Dr. E.H.C. McKenzie, Dr D.W. Li, and Prof. Y.L. Guo. This project was supported by the National Science Foundation of China (no. 31060005).

Literature cited

- Ellis MB. 1971. Dematiaceous hyphomycetes. X. Mycological Papers 125: 1–30.
- Hansford CG. 1943. Contributions towards the fungus flora of Uganda. V. Fungi imperfecti. Proceedings of the Linnean Society of London 155: 34–67. doi: 10.1111/j.1095-8312.1943.tb00340.x
- Jiang YL, Zhang TY. 2007. Notes on soil dematiaceous hyphomycetes from Shennongjia Natural Conservation Area, Hubei Province I. Mycosystema 26(1): 17–21. doi: CNKI:SUN:JWXT.0.2007-01-004
- Karandikar KG, Kulkarni SM, Patwardhan PG. 1992. Some new and interesting hyphomycetes from India. Biovigyanam 18(2): 78–81.
- Matsushima T. 1971. Mycological reports from New Guinea and the Solomon Islands. 7. Some interesting fungi imperfecti. Bulletin of the National Science Museum, Tokyo, 14: 460–480.
- McKenzie EHC. 1991. Dematiaceous hyphomycetes on *Freycinetia* (*Pandanaceae*). 1. *Stachybotrys*. Mycotaxon 41: 179–188.
- Pinruan U, McKenzie EHC, Jones EBG, Hyde KD. 2004. Two new species of *Stachybotrys*, and a key to the genus. Fungal Diversity 17: 145–157.
- Tubaki K. 1963. Notes on the Japanese hyphomycetes. I. *Chloridium*, *Clonostachys*, *Ishmospora*, *Pseudobotrytis*, *Stachybotrys* and *Stephanoma*. Transactions of the Mycological Society of Japan 4: 83–90.
- Whitton SR, McKenzie EHC, Hyde KD. 2001. Microfungi on the *Pandanaceae*: *Stachybotrys*, with three new species. New Zealand Journal of Botany 39: 489–499. doi: 10.1080/0028825X.2001.9512752